

Bryophyte flora of the Huon Peninsula, Papua New Guinea. LXXII. Spiridentaceae (MUSCI)

Daniel H. Norris¹, Timo Koponen² & J. R. Shevock¹

¹⁾ University Herbarium, University of California, Berkeley, CA 94720, U.S.A

²⁾ Timo Koponen, Botanical Garden and Museum (Bryology), P.O. Box 7, FI-00014 University of Helsinki, Finland, and: Finnish-Chinese Botanical Foundation, Mailantie 109, FI-08800 Lohja, Finland; e-mail: timo.koponen@helsinki.fi

Received April 20, 2010, accepted Oct. 17, 2010

Norris, D. H., Koponen, T. & Shevock, J. R. 2010: Bryophyte flora of the Huon Peninsula, Papua New Guinea. LXXII. Spiridentaceae (Musci). — In: Koponen, T., Piippo, S. & Reinikka, E. (eds.), Dr. Ming-Jou Lai Memorial Volume. — *Acta Bryolichenologica Asiatica* 3: 157–171.

In Western Melanesia *Spiridens* Nees has 3 species: *S. longifolius* Lindb., *S. perichaetialis* E. B. Bartram & Willis in E. B. Bartram, and *S. reinwardtii* Nees. *Spiridens armatus* E. B. Bartram and *S. balfouriana* Greville are synonymized with *S. reinwardtii*. *S. flagellosus* Mitt. and *S. aristifolius* are excluded from the flora. The paper includes keys to the species, descriptions, illustrations and distributional data for the species in Western Melanesia. The family Spiridentaceae and the genus *Spiridens* are described and each species description includes discussions on taxonomy and classification, as well as nomenclatural notes and range descriptions.

Key words: distribution, Indonesia, nomenclature, Papua New Guinea, Solomon Islands, *Spiridens*, Spiridentaceae, taxonomy, West Irian